PEDIATRIC INTERFACILITY TRANSFER FORM

| DATE: | | | AGE: | | | | MEDICAL RECORD NUMBER # | | | | | | | | | | |
|--|--------|----------------|------------------|--|----------|--------------|-------------------------|------|---------|--------------|-----------------|-----------------|------------------|------------|-------------|-------|---|
| DIAGNOSIS: | | | | | | | | | | | | | | | | | |
| ARRIVAL MODE TO THE ED: PRIVATE VEHICLE AMBULA | | | | | ANCE | NCE | | | | | | | | | | | |
| ASSESSMENT PRIOR 1 | ГО ТНЕ | TRANSF | ER: | | | | | | | | | | | | | | |
| CNS RES | | | PIRATORY STATUS | | | | CARDIOVASCULA | | | R | | MUSCULOSKELETAL | | | VITAL SIGNS | | |
| ALERT | Y/N | INTUBATED Y/ | | | Y/N | I/V | | | | Y/N | CERVICAL COLLAR | | Y/N/NA | B/P | | | |
| CONSCIOUS Y/N TRAC | | | CHEOTOMY | | | Y/N | I/O | | | | Y/N | BAC | BACK BOARD Y/N/N | | PULSE: | | |
| GLASGOW COMA SCALE () | | CRICOTHYROTOMY | | | Y/N | CENTRAL LINE | | | Y/N | | EXTREMITY: | | | RESP RATE: | | | |
| | | PULSE OXIMETER | | | Y/N | | | | | | | | | | | | |
| A V P U OXY | | | GEN:L BVM/NC/MAS | | | ASK | OTHER: | | | | | OTHER: | | | TEMP: | | |
| MODE OF TRANSPORT | : AM | BULANC | E | | AIR TRAN | SPORT | | 0 | THER (S | PECIFY) | | | | | • | | |
| LEVEL OF PROVIDER TRANSPORT: BLS | | | | | ILS | | ALS | LS / | | | ACCO | MPANIED BY: MD | | R | N | OTHER |] |
| RECEIVING FACILITY: | | | | | | | | | | | | | | | | | |
| RECEIVING MD: | | | | | | | | | | SENDING MD: | | | | | | | |
| REASON FOR TRANSFER: 1. NEED FOR HIGHER LEVEL OF CARE 2. NEED FOR SPECIALTY CARE 3. FAMILY/MD REQUEST 4. INSURANCE | | | | | | | 5. OTHER (EXPLAIN) | | | | | | | | | | |
| | | | | | | | YES | NO | N/A | N/A COMMENTS | | | | | | | |
| PRIVATE MD NOTIFIED? | | | | | | | | | | | | | | | | | |
| CONSULTATION WITH TERTIARY CARE CENTER? | | | | | | | | | | | | | | | | | |
| TRANSFER SHEET DOCUMENTATION COMPLETED? | | | | | | | | | | | | | | | | | |
| IS TOTAL TIME IN ED < 3 HOURS? IF NOT, PLEASE COMMENT. | | | | | | | | | | | | | | | | | |
| CONDITION OF PATIENT AT TIME OF TRANSFER DOCUMENTED? | | | | | | | | | | | | | | | | | |
| Patient belongings given to family member for transfer to receiving institution. | | | | | | | | | | | | | | | | | |
| Famil given directions to recieiving hospital and accepting MD name and contact information | | | | | | | | | | | | | | | | | |

INSTRUCTIONS FOR COMPLETING THE PEDIATRIC INTERFACILITY TRANSFER FORM

CENTRAL NERVOUS SYSTEM ASSESSMENT:

- Glasgow Coma Scale see attachment
- A V P U is a level of consciousness assessment
 - A = Alert
 - V = Responds to verbal stimulus
 - P = Responds to painful stimulus
 - U = Unresponsive to any stimulus

RESPIRATORY STATUS:

- Begin with assessing if respiratory system is stable or not.
- If not, answer yes or no to the four questions. Indicate how much oxygen is being administered and whether by bag valve and mask (BVM), nasal cannula (NC), or mask

CARDIO-VASCULAR ASSESSMENT:

Indicate vascular access as either: intravenous, intraosseous, central line or other.

VITAL SIGNS:

Last vitals before transfer

DESIGNATED CHILD LIAISON:

A specific person who is designated to stay with the child throughout the entire emergency room stay, including moves to radiology etc. Some hospitals designate a respiratory therapist, a chaplain, a social worker. Indicate whether someone met this role and what their background was.

TOTAL TIME IN ED:

Time from arrival until transfer departure

MODIFIED GLASCOW COMA SCALE

FOR INFANTS/CHILDREN

ACTIVITY

BEST RESPONSE

| Eye Opening: | |
|--|-----------------------|
| SpontaneousTo SpeechTo PainNone | 4 3 2 1 |
| Verbal: | |
| Coos, Babbles. Irritable Cries to Pain Moans or Grunts to Pain None | 5 4 3 2 1 |
| Motor: | |
| Normal Spontaneous movements. Withdraws to Touch. Withdraws to Pain. Abnormal Flexion (decorticate). Abnormal Extension (decerebrate). | 6 5 4 3 2 |

GLASCOW COMA SCALE

| EYE RESPONSE | | | |
|---------------------------|---------------------------|----------------|---------------|
| Automatic (opens eyes | | | |
| To voice (opens eyes w | hen told to do so) | | 3 |
| To pain (opens eyes on | ly to painful stimulus) | | 2 |
| None (does not open ey | es to any stimulus) | | 1 |
| VERBAL RESPONSE (Asked "V | | | |
| Confused (tells you cor | | | |
| Inappropriate reply (ran | dom renty) | ,t answer j | |
| Unintelligent sounds (in | comprehensible moans | s) | 2 |
| None (no verbal respon | | | |
| ` . | , | | |
| MOTOR RESPONSE | | | _ |
| Obeys verbal direction | (shows two fingers whe | n asked) | 6 |
| Controlled movements | (points to pain location) | | 5 |
| Withdraws (moves away | | | |
| Abnormal flexion (assu | mes a decorticate postu | re) | 3 |
| | | | |
| | | _ | |
| | | 7 | |
| | | THE ST | |
| - | | | |
| | | ماجر | |
| | | | |
| Abnormal sytopoion (so | oumas a dagarahrata na | noturo) | 2 |
| Abnormal extension (as | ssumes a decerebrate po | osture) | 2 |
| | | | |
| | £ | | |
| | | \sim | |
| | | | |
| | |) [2] | |
| | | <u></u> .) | |
| | | | |
| | | | |
| None (no response, lies | flaccid) | | l |
| | | T0T41 | |
| | | TOTAL POINTS _ | |
| Glasgow Coma Scale Range | | | |
| | | | |
| | 12 9 | 6 3 | |
| Normal | ?Neurological damage | 9 | Severe damage |